SCIENTIFIC REPORTS

Received: 21 July 2015 Accepted: 28 September 2015 Published: 23 October 2015

OPEN Anaerobic ammonium oxidation and its contribution to nitrogen removal in China's coastal wetlands

Lijun Hou^{1,*}, Yanling Zheng^{1,2,*}, Min Liu^{2,*}, Xiaofei Li², Xianbiao Lin², Guoyu Yin^{1,2}, Juan Gao¹, Fengyu Deng¹, Fei Chen¹ & Xiaofen Jiang¹

Over the past several decades, human activities have caused substantial enrichment of reactive nitrogen in China's coastal wetlands. Although anaerobic ammonium oxidation (anammox), the process of oxidizing ammonium into dinitrogen gas through the reduction of nitrite, is identified as an important process for removing reactive nitrogen, little is known about the dynamics of anammox and its contribution to nitrogen removal in nitrogen-enriched environments. Here, we examine potential rates of anammox and associate them with bacterial diversity and abundance across the coastal wetlands of China using molecular and isotope tracing techniques. High anammox bacterial diversity was detected in China's coastal wetlands and included Candidatus Scalindua, Kuenenia, Brocadia, and Jettenia. Potential anammox rates were more closely associated with the abundance of anammox bacteria than to their diversity. Among all measured environmental variables, temperature was a key environmental factor, causing a latitudinal distribution of the anammox bacterial community composition, biodiversity and activity along the coastal wetlands of China. Based on nitrogen isotope tracing experiments, anammox was estimated to account for approximately 3.8-10.7% of the total reactive nitrogen removal in the study area. Combined with denitrification, anammox can remove 20.7% of the total external terrigenous inorganic nitrogen annually transported into China's coastal wetland ecosystems.

Reactive nitrogen (Nr) production has increased over the past several decades^{1,2}. Global Nr enrichment is one of the main emerging environmental problems of the twenty-first century³⁻⁵. Nr is now entering Earth's land surface at more than twice its natural rate due to industrial production of nitrogen fertilizer and combustion of fossil fuels^{2,4}. On a global basis, approximately 160 teragrams (Tg) of anthropogenic nitrogen are added to the Earth's surface ecosystem each year⁴ in efforts to sustain food production for an increasing population and as parts of other social and economic human activities⁶. Approximately 20-30% of the anthropogenic nitrogen produced during fossil fuel combustion and food production and consumption is transported to estuarine and coastal marine systems via river flow^{3,7}. This Nr pollution presents a serious threat to the environmental quality and health of these ecosystems in many areas of the world^{8,9}, as increasing nitrogen input into estuarine and coastal marine systems is an important driver of water pollution (e.g., coastal eutrophication, hypoxia, harmful algae blooms, and seawater acidification)9,10.

China is seriously affected by Nr pollution^{7,11}. Over the past century, Nr production in mainland China has increased from approximately 9.2 to 56 Tg per year¹¹, with an annual growth rate of 1.8%.

¹State Key Laboratory of Estuarine and Coastal Research, East China Normal University, Shanghai 200062, China. ²College of Geographical Sciences, East China Normal University, Shanghai 200241, China. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to L.J.H. (email: ljhou@sklec.ecnu.edu.cn) or M.L. (email: mliu@geo.ecnu.edu.cn)





Since the 1950s, anthropogenic Nr creation from agriculture-associated biological nitrogen fixation, industrial nitrogen fixation, and fossil fuel combustion has exceeded natural Nr creation from lightning and biological nitrogen fixation in forests and grasslands¹¹. Over the past three decades (1980–2010), anthropogenic nitrogen production was exponentially linked to gross national production, as well as to gross agricultural and industrial production (Supplementary Fig. S1). These strong associations imply that the input of anthropogenic nitrogen has greatly contributed to food and energy production in China. However, a large fraction of this human-produced nitrogen is delivered to atmospheric and aquatic environments¹² due to low nitrogen-use efficiency by fertilized crops and other organisms. Nearly 20% of the total lost Nr, mostly in the form of nitrate, is transported into estuarine and coastal ecosystems by riverine discharge and atmospheric deposition¹¹. As a result, the estuaries and coastal seas of China have become highly Nr-enriched ecosystems, characterized by extreme hyper-eutrophication and algal bloom formation (Supplementary Fig. S2). An improved understanding of nitrogen transformation and fate on a national scale is required to develop strategies to protect the water quality and health of these estuarine and coastal environments.

Anaerobic ammonium oxidation (anammox), the process of oxidizing ammonium through the reduction of nitrite, removes Nr permanently from aquatic ecosystems and returns it to the atmosphere as nitrogen gas¹³, as does denitrification. Anammox was first discovered in a wastewater treatment plant (WWTP) in the Netherlands in 1995¹⁴. Since that discovery, this process has been detected in various natural ecosystems, including freshwater sediments, marine sediments, and anaerobic water columns^{13,15-21}. The anammox process has also been observed in the coastal wetlands of China^{21,22}. However, these studies focused on local and small-scale regions. Gaps remain in our knowledge of the anammox process, including its association with anammox bacterial dynamics and its contribution to nitrogen loss across the coastal wetlands of China. To fill in these knowledge gaps, sediments (0–5 cm deep) were collected from the coastal wetlands of China in January and in August 2014 (Fig. 1), and molecular and isotope tracing experiments were conducted to obtain data on anammox bacteria and activity. This work examines for the first time the latitudinal distributions of anammox bacteria with regard to community composition, biodiversity, and activity, providing novel insights into the microbial nitrogen cycle in coastal wetlands.

Results

Anammox Rates. Anammox rates were measured at all sampling sites along the coastal wetlands of China using sediment slurry incubation experiments combined with a ¹⁵N isotope-labeling technique^{23,24}. We conducted three independent groups of sediment slurry incubation experiments to quantify the rates of anammox and its contribution to N₂ production, spiking each group with ¹⁵NH₄⁺, ¹⁵NH₄⁺ + ¹⁴NO₃⁻, or ¹⁵NO₃⁻. No significant production of isotope-labeled dinitrogen gas, including ²⁹N₂ and ³⁰N₂, was detected from the incubations with ¹⁵NH₄⁺ alone in any of the collected sediment samples (Supplementary Fig. S3a), indicating complete consumption of background nitrate and nitrite in the sediment slurries during pre-incubation. With the addition of ¹⁵NH₄⁺ + ¹⁴NO₃⁻, significant production of ²⁹N₂ was detected in all sediment samples, but no significant production of ³⁰N₂ was observed (Supplementary Fig. S3b). The results from these two groups of experiments corroborated the occurrence



Figure 2. Spatiotemporal variations of anaerobic ammonium oxidation (anammox) rates (a) and associated contribution (b) to nitrogen loss at each sampling site. Vertical bars indicate standard error (n = 3).

of anammox at all sampling sites. By contrast, significant production of both $^{29}N_2$ and $^{30}N_2$ was detected from incubations containing added $^{15}NO_3^-$ (Supplementary Fig. S3c). Therefore, rates of anammox and its contributions to N_2 production were quantified from the experiments in which $^{15}NO_3^-$ was added 15,25 .

Potential rates of anammox obtained in the sediment slurry incubation experiments ranged from 1.8 to 10.4μ mol N kg⁻¹ d⁻¹ (Fig. 2a). Anammox rates were significantly greater in summer than in winter (one-way ANOVA, p < 0.05), with respective average values of $7.0 \pm 1.4 \mu$ mol N kg⁻¹ d⁻¹ and $3.2 \pm 0.8 \mu$ mol N kg⁻¹ d⁻¹ (Supplementary Fig. S4). Significantly lower rates of anammox were recorded at coastal wetland sites S1 to S6 than at sites S7 to S11 (one-way ANOVA, p < 0.05), with average values of $5.9 \pm 2.3 \mu$ mol N kg⁻¹ d⁻¹ and $4.3 \pm 1.9 \mu$ mol N kg⁻¹ d⁻¹, respectively (Supplementary Fig. S5). Among all measured environmental variables (Supplementary Table S1), winter anammox rates were significantly related (Pearson's correlation analyses) to temperature (R = 0.752, p = 0.008), nitrite (R = 0.667, p = 0.024), and ammonium (R = 0.833, p = 0.002), whereas summer anammox rates were significantly correlated with temperature (R = 0.607, p = 0.048), nitrate (R = 0.649, p = 0.030), nitrite (R = 0.692, p = 0.018), and organic carbon (R = 0.717, p = 0.012) (Supplementary Table S2). The estimated contributions of anammox to total N₂ production varied from 3.8 to 10.7% in winter and from 4.5 to 5.8% in summer (Fig. 2b); the remainder would be attributed to denitrification.

Composition, Diversity, and Abundance of Anammox Bacteria. To further decipher the microbial mechanisms regulating anammox, the community composition, diversity, and abundance were examined based on the 16S rRNA gene of anammox bacteria. Phylogenetic analysis of the 16S rRNA genes indicated the presence of four known anammox bacterial genera, *Candidatus Scalindua, Kuenenia, Brocadia,* and *Jettenia,* as well as a novel cluster (Supplementary Fig. S6). Weighted UniFrac PCoA analysis with the 16S rRNA gene sequences showed no significant seasonal shift in the community composition or diversity of the anammox bacteria at most of the sampling sites (except for sites S2 and S4; Supplementary Fig. S7). However, a distinctive latitudinal differentiation was observed in the community composition and diversity of the anammox bacteria along the coastal wetlands of China (Fig. 3). The community of anammox bacteria at sites S1 to S6 was mainly dominated by *Scalindua*, which accounted for >90% of the sequences (Supplementary Fig. S8). By contrast, significant shifts were recorded in the compositions of the anammox bacteria communities at sites S7 to S11, which consisted mainly of *Scalindua*, *Kuenenia*, and (or) *Brocadia*. Potential relationships between these anammox bacterial distributions and



Figure 3. UniFrac weighted PCoA analysis of anaerobic ammonium oxidation (anammox) bacterial assemblages as revealed by the 16S rRNA gene sequences. S and W in parentheses represent summer and winter samples, respectively.





environmental factors were examined by weighted canonical correspondence analysis (CCA). Of all the environmental factors investigated, only temperature had a significant effect on the anammox bacterial community distribution (p = 0.002 for winter; p = 0.038 for summer; Supplementary Fig. S9).

The abundance of anammox bacteria in the coastal wetland sediments of China was estimated with qPCR assays based on the 16S rRNA gene. The quantity of anammox bacterial 16S rRNA genes observed ranged between 1.17×10^7 and 4.25×10^7 copies g⁻¹ dry sediment (Fig. 4). Anammox bacterial abundance was characterized by significant spatiotemporal heterogeneity at the study area (one-way ANOVA, p < 0.05). The abundance of anammox bacteria was generally higher in summer than in winter, with average abundances of $2.87 \pm 0.8 \times 10^7$ copies g⁻¹ dry sediment in summer and $2.05 \pm 0.6 \times 10^7$ copies g⁻¹ dry sediment in winter (Supplementary Fig. S10). Additionally, the spatial distribution of anammox bacterial abundance was characterized by an increasing trend from high to low latitudes (Supplementary Fig. S11). Pearson's correlation analyses revealed that the anammox bacterial abundance was related significantly to temperature (R=0.862, p < 0.0001) and the amounts of sedimentary ammonium (R=0.814, p=0.002) and nitrite (R=0.795, p=0.006) in winter, whereas it was closely related to temperature (R=0.790, p=0.004) and the amounts of sedimentary nitrate (R=0.802, p=0.004) and nitrite (R=0.856, p < 0.0001) in summer (Supplementary Table S3).

Other Nitrogen Transformation Processes. Potential rates of related nitrogen cycling processes, including denitrification, nitrification and dissimilatory nitrate reduction to ammonium (DNRA), were





.....

determined to examine their potential links to anammox (Fig. 5). Denitrification rates in sediments in the study area ranged from 20.8 to 78.9 µmol N kg⁻¹ d⁻¹ in winter. Significantly higher denitrification rates were observed in summer (one-way ANOVA, p < 0.05), with values of 87.8–182.1 µmol N kg⁻¹ d⁻¹. In winter, potential nitrification rates in sediments varied between 10.9 and 19.6 µmol N kg⁻¹ d⁻¹. However, the relatively higher rates of nitrification in summer ranged from 12.9 to 29.3 µmol N kg⁻¹ d⁻¹ (one-way ANOVA, p < 0.05). Compared to denitrification and nitrification, DNRA showed relatively low rates, with values of 3.5–10.1 µmol N kg⁻¹ d⁻¹ in winter and 11.4–19.5 µmol N kg⁻¹ d⁻¹ in summer. Pearson's correlation analyses showed that summer anammox rates were significantly correlated with denitrification rates (R = 0.870, p = 0.0004), whereas in winter, anammox rates correlated with nitrification rates (R = 0.745, p = 0.008) (Fig. 6). Anammox and DNRA rates did not correlate significantly in the study area (p > 0.05).

Discussion

Both molecular and isotope-tracing experiments corroborated the occurrence of anammox in the coastal wetland sediments of China. The anammox process is mediated by chemolithoautotrophic bacteria affiliated with the order *Brocadiales*, part of the phylum Planctomycetes²⁶. At present, five genera of anammox bacteria (*Candidatus Brocadia, Kuenenia, Scalindua, Anammoxoglobus* and *Jettenia*) have been identified from different wastewater treatment systems²⁶. Using molecular techniques, the biodiversity



Figure 6. Pearson's correlations of anaerobic ammonium oxidation (anammox) rates with denitrification, nitrification and dissimilatory nitrate reduction to ammonium (DNRA) rates in China's coastal wetlands. Vertical and horizontal bars indicate standard error (n=3).

.....

of anammox bacteria was recently studied in various natural ecosystems and determined to be relatively low for marine, riverine, and lacustrine ecosystems²⁷⁻³⁰; community compositions in these environments were restricted mainly to Scalindua or Brocadia²⁷⁻³⁰. By contrast, a relatively high biodiversity of anammox bacteria, including Candidatus Scalindua, Kuenenia, Brocadia, and Jettenia (based on the 16S rRNA gene), was detected in our study area. These results are similar to the high biodiversity of anammox bacteria observed in other estuarine and coastal wetlands^{19,31,32}. Therefore, this comparison implies that high anammox bacterial biodiversity may occur in the transitional zones between terrestrial and marines ecosystems, where diverse biotopes and eco-niches exist in complex ecotones. Although the biodiversity of anammox bacteria did not show a statistically significant seasonal shift, it was characterized by a distinctive latitudinal differentiation along the coastal wetlands of China. Numerous studies have reported that salinity is a key environmental factor shaping the diversity and biogeographical distribution of anammox bacteria^{21,27,28,33}. However, in our study area, the distribution of the anammox bacterial community was not significantly related to salinity, but it did correlate with temperature (Supplementary Fig. S9). Compared with the high latitude sites, low latitude sites contained more diverse communities of anammox bacteria. These results imply that temperature is a key environmental factor shaping the biogeographical distribution and diversity of the anammox bacterial community in the coastal wetlands of China. Therefore, differences in the eco-physiology of these anammox bacteria might be a main mechanism defining the geographical distribution pattern of the anammox bacterial community. The latitudinal distribution pattern observed in the present study emphasizes the importance of temperature in controlling the biogeographical distribution of anammox bacterial community structure and diversity at a larger spatial scale.

Anammox bacterial abundance demonstrated significant spatial and seasonal fluctuations in the study area. Similarly, temperature was a key environmental variable controlling the spatio-temporal heterogeneity of anammox bacteria abundance in sediments (Pearson's correlation analyses; Supplementary Table S3). The significant correlation between nitrite and anammox bacteria abundance implies that substrate availability is also an important factor affecting the number of anammox bacteria in coastal wetland sediments^{15,34,35}. Nitrite may help limit anammox bacterial population development in the coastal wetland sediments of China. However, a seasonal shift was observed in the underlying mechanism controlling substrate availability. In winter, nitrite may derive from nitrification. Significant associations between sedimentary nitrite content and nitrification rate (R=0.609, p=0.047) as well as sedimentary ammonium content (R=0.710, p=0.014) support this conclusion. By contrast, nitrite is more likely produced from denitrification rate (R=0.700, p=0.016) and sedimentary nitrate content (R=0.829, p=0.002).

Anammox rates and contribution to nitrogen removal were examined in the coastal wetland sediments of China. Our isotopic tracing experiments revealed rates of anammox that are comparable to the values reported in other estuarine and coastal wetland sediments (Table 1). Anammox rates in the coastal wetland sediments of China were characterized by substantial spatial and seasonal heterogeneity,

| Locations | Rates (µmol N m ⁻² h ⁻¹) | Contributions (%) | Refs |
|-----------------------------------|---|-------------------|------------|
| New River Estuary | 0.3-50.8ª | <14 | 19 |
| Seine Estuary | 11.8-24.9ª | 3-8 | 20 |
| Rander Fjord | 19–55 | 5-24 | 23 |
| Norsminder Fjord | | | |
| Thames Estuary | 2.2-200 | 0.5-8 | 25 |
| Cape Fear River Estuary | 1.5-14.8 ^a | 3.8-14.6 | 31 |
| Southeast coast of Queensland | 0-84 | 0-9 | 34 |
| West and south coast of Greenland | 1–92 | 1-35 | 42 |
| Gulf of Finland | 10-30 | 10-15 | 43 |
| Chesapeake Bay | 0-1.6 | 0-22 | 44 |
| Colne Estuary | 0-157 | <30 | 45 |
| Plum Island Sound Estuary | 2.9 | <3 | 46 |
| St. Lawrence Estuary | 5.5 | 3 | 47 |
| Providence River Estuary | 0-16 | 0-4 | 48 |
| Little Lagoon | 0.2-0.5 | 2-4 | 49 |
| China's Coastal wetlands | 2.0-11.4ª | 3.8-10.7 | This study |

Table 1. Rates of anaerobic ammonium oxidation (anammox) and associated contribution to nitrogen removal in China's coastal wetlands and other estuarine and coastal ecosystems. ^aThe unit is changed from μ mol N g⁻¹h⁻¹ to μ mol N m⁻²h⁻¹, assuming dry sediment bulk density of 1.88 g cm⁻³.

.....

which might relate to sampling site biotic and abiotic factors. In the present study, anammox rates were not significantly related to the anammox bacterial community composition and structure (p > 0.05; Supplementary Fig. S12) but did correlate with their abundance (R = 0.852, p = 0.0008 for winter; R = 0.633, p = 0.036 for summer; Supplementary Fig. S13). This result demonstrates that anammox bacterial abundance, more than composition and diversity, predicts anammox activity³¹. In addition, anammox rates were significantly and positively correlated with temperature (R = 0.752, p = 0.008 for winter; R = 0.607, p = 0.048 for summer). High temperature may favor the metabolism of anammox bacteria and thus promote rates of anammox^{22,36}. The relationships between anammox rates and sedimentary nitrite concentrations (R = 0.667, p = 0.024 for winter; R = 0.692, p = 0.018 for summer) imply that substrate availability is a limiting factor affecting the activity of anammox bacteria in the coastal wetland sediments of China.

Anammox rates were compared to nitrification, denitrification and DNRA rates to elucidate the potential links among these nitrogen transformation processes. Anammox rates were not significantly related to DNRA rates but did relate to nitrification and denitrification rates (Fig. 6). These correlation analyses reflected that anammox might be closely associated with nitrification and denitrification in the study area. Interestingly, the links between anammox and nitrification and denitrification shifted seasonally. In winter, anammox correlated predominantly with nitrification, suggesting partial nitrification as a primary source of nitrite for winter anammox reactions. This conclusion is also supported by the significant relationships of anammox rates with sedimentary ammonium (R = 0.833, p = 0.002) and nitrite (R = 0.667, p = 0.024) contents. By contrast, summer anammox shifted to coupling with denitrification. During the interaction between anammox bacteria and denitrifiers, denitrification was likely a primary source of nitrite for anammox in the coastal wetland sediments of China³⁴. The strong linkage of both processes is supported by the significant correlations of anammox rates with sedimentary nitrate (R = 0.649, p = 0.030) and organic carbon (R = 0.717, p = 0.012) contents. In summer, more nitrite was generated as an intermediate product by enhanced denitrification (Supplementary Fig. S14). As a result, the development of anammox bacteria was stimulated (Supplementary Table S3) and enhanced anammox rates in the warm season (Supplementary Fig. S13b). Although the DNRA process can also produce nitrite as an intermediate product³⁶ and thereby act as an alternative source of substrate for anammox bacteria^{16,18}, no significant relationship was detected between anammox and DNRA rates (p > 0.05), implying that DNRA-produced nitrite contributes minimally to the metabolic activity of anammox bacteria in nitrogen-enriched coastal wetlands where denitrification and nitrification are active³⁷.

We estimated that the anammox process, on average, contributed approximately $5.7 \pm 1.2\%$ to total nitrogen loss from the coastal wetland sediments of China; this approximation is comparable to that reported in rivers³⁸, lakes^{39,40}, constructed wetlands⁴¹, and other estuarine and coastal environments^{23,24,42-49} but lower than that generally observed in marine ecosystems¹⁵⁻¹⁸. If the average rate of anammox is extrapolated to the entire study area (5.8×10^6 ha), approximately $2.81 \pm 1.1 \times 10^4$ t N can be removed annually from the coastal wetland ecosystem by this process, based on dry sediment bulk

density (1.88 g cm⁻³) in the study area (Supplementary Fig. S15). This removal accounts for approximately $1.2 \pm 0.5\%$ of the total multi-annual average terrigenous inorganic nitrogen (2.4×10^6 t N) transported into the coastal wetlands of China. If denitrification is also included, both processes can remove approximately $4.98 \pm 1.87 \times 10^5$ t N from the study area on an annual basis, which accounts for approximately $20.7 \pm 7.8\%$ of the total external terrigenous inorganic nitrogen. Furthermore, assuming that the coastal wetlands of China have not been destroyed by coastal development in recent decades, up to $49.3 \pm 18.6\%$ of terrigenous inorganic nitrogen could be removed by both denitrification and anammox, which would greatly benefit the ecological health of China's coastal seas.

Taken as a whole, this study is the first to investigate anammox rates, the contribution of anammox rates to nitrogen removal, and the dynamics of anammox bacteria in the coastal wetlands of China on a national scale. High biodiversity of anammox bacteria was observed in the coastal wetlands and included *Candidatus Scalindua, Kuenenia, Brocadia*, and *Jettenia*. Compared to denitrification, anammox contributed approximately $5.7 \pm 1.2\%$ of the total loss of reactive nitrogen at the study area. The present study highlights the importance of temperature in regulating the latitudinal distribution of the anammox bacterial community composition, diversity, abundance, and activity in natural environments on a large spatial scale. This study also suggests the significance of protecting coastal wetlands for the sustainable development of society and the economy in China's coastal zone.

Methods

Study Area. The Chinese coastal zone covers an area of more than three million square kilometers with an 1800-km coastline stretching across tropical, subtropical and temperate zones⁵⁰. Coastal development in China has contributed greatly to the national economy. The gross domestic productivity (GDP) in the coastal regions accounted for 50% of China's total GDP before 1978 and increased to 60% between 1978 and 2010^{51} . Meanwhile, China's coastal population increased from 260 million in 1954, to 400 million in 1978, and 590 million in 2010^{51} . The continuing increase in population coupled with economic growth, rapid urbanization, and infrastructure development in the coastal zone of China's coastal wetlands have suffered tremendous loss due to increased threats and pressures on wetlands by the increasing population and rapid economy growth. From 1950 to 2014, the area of coastal wetlands decreased from 13.81×10^6 ha to 5.8×10^6 ha, with a total loss rate of $58\%^{52}$. Numerous factors endanger the existence of coastal wetlands, such as land demands by a large human population, a lack of understanding of coastal wetland values, a misguided reclamation policy, and a lack of environmental laws and regulations. Among them, reclamation and infrastructure construction were the primary causes of wetland loss, accounting for 70-82% of the total loss⁵³.

Sample Collection. In this study, eleven sampling sites were selected along the coastal wetlands of China (Fig. 1). Field surveys were carried out in January and in August 2014. Triplicate sediment samples (0–5 cm deep) were collected from each site with PVC tubes. Sediment cores were stored in sterile plastic bags, sealed and transported to the laboratory on ice within 4–8 h. The sediment of each core was mixed thoroughly under helium to form one composite sample. One portion of the homogenized sediment was incubated immediately to measure nitrogen transformation rates, and another portion was examined for sediment physiochemical characteristics. Subsamples were also preserved at -80 °C for DNA extraction and molecular analysis.

Analysis of Sediment Characteristics. Sediment temperature was measured *in situ* with a portable electronic thermometer at each site. Sediment salinity and pH were measured using a YSI Model 30 salinity meter and a Mettler-Toledo pH meter, respectively, after the sediments were mixed with deionized water free of CO₂ at a sediment/water volume ratio of $1:2.5^{22}$. Sediment grain size was measured using a LS 13 320 Laser grain sizer. Organic carbon (OC) was determined by a CHN elementary analyzer (VVarioELIII, Elementary, Germany) after removing carbonate by leaching with 0.1 mol L⁻¹ HCl²². Exchangeable ammonium (NH₄⁺-N), nitrite (NO₂⁻-N), and nitrate (NO₃⁻-N) were extracted from fresh sediments with 2 mol L⁻¹ KCl and measured spectrophotometrically on a continuous-flow nutrient analyzer (SAN plus, *Skalar Analytical* B.V., the Netherlands), with detection limits of 0.5 µmol L⁻¹ for NH₄⁺-N and 0.1 µmol L⁻¹ for NO₂⁻-N and NO₃⁻-N²². Sulfide was measured using an Orion Sure-flow[®] combination silver-sulfide electrode (Thermo Scientific Orion)⁵⁴, with a detection limit of 0.09 µmol L⁻¹. All sediment physiochemical parameters were analyzed in triplicate. Detailed information on sediment characteristics is given in Supplementary Table S1.

DNA Isolation, PCR, Cloning, Sequencing, and Phylogenetic Analysis. Total genomic DNA was extracted from 0.25g of each sediment sample using PowersoilTM DNA Isolation Kits (MOBIO, USA) according to the manufacturer's protocol. Nested PCR assays were conducted to detect the anammox bacterial 16S rRNA gene. The initial PCR reaction was conducted using PLA46f-1390r primers with a thermocycling program of 94 °C for 4 min followed by 30 cycles of 45 s at 95 °C, 50 s at 59 °C, 1 min 40 s at 72 °C, and a final 5 min extension cycle at 72 °C²². One μ L of the 10-fold diluted PCR product was then used as a template for the second amplification with Amx368f-Amx820r primers using a thermocycling

program of 94 °C for 4 min followed by 30 cycles of 45 s at 95 °C, 50 s at 59 °C, 1 min at 72 °C, and a final 5 min extension cycle at 72 °C²².

Appropriately sized fragments of the nested PCR products were separated by electrophoresis in 1% agarose gels and purified using the Gel Advance-Gel Extraction system (Viogene, China). The purified fragments were cloned by TOPO-TA cloning (Invitrogen, USA) in accordance with the manufacturer's instructions. Clones were selected randomly for further analysis. The unique sequences obtained in this study for anammox bacteria are available in NCBI under accession numbers KT229750-KT230483. All of the sequences and their relatives obtained from NCBI were aligned using the ClustalX program (version 2.1)⁵⁵. Sequences with 97% identity were grouped into one operational taxonomic unit (OTU) using the Mothur program (version 1.33.0, USA) (http://www.mothur.org/wiki/Main_Page) by the furthest neighbor approach. Phylogenetic trees were constructed by the neighbor-joining (NJ) method using MEGA software (version 5.03)⁵⁶. The relative confidence of the tree topologies was evaluated by performing 1000 bootstrap replicates⁵⁷.

Real-Time Quantitative PCR Assay. Plasmids carrying an anammox bacterial 16S rRNA gene fragment were extracted from *Escherichia coli* hosts using a Plasmid Mini Preparation Kit (Tiangen, China) for standard curve construction. Plasmid DNA concentrations were measured with a Nanodrop-2000 Spectrophotometer (Thermo, USA). Triplicate samples and standard reactions were performed with primers AMX-808-F and AMX-1040-R on an ABI 7500 Sequence Detection System (Applied Biosystems, Canada)²². The 25µL qPCR mixture contained 12.5µL of Maxima SYBR Green/RoxqPCR Master Mix (Fermentas, Lithuania), 1µL of each primer (10µmol L⁻¹) and 1µL template DNA. All reactions were performed in 8-strip thin-well PCR tubes with ultraclear cap strips (ABgene, United Kingdom). PCR amplification was performed as follows: 50 °C for 2 min, 95 °C for 10 min, followed by 45 cycles of 30 s at 95 °C, 30 s at 55 °C, 30 s at 72 °C. The specificity of the Q-PCR amplification was determined by melting curve and gel electrophoresis. In all experiments, negative controls containing no template DNA were subjected to the same qPCR procedure to detect and exclude any possible contamination or carryover.

Melting-curve analyses of the anammox bacterial 16S rRNA revealed a single peak at 84.3 °C, corroborating that fluorescent signals were derived from specific PCR products in the process of qPCR quantifications. Standard curves were generated from plasmids containing cloned 16S rRNA gene fragments by plotting the threshold cycle (Ct) versus the \log_{10} value of the gene copy number. A significant linear relationship ($R^2 = 0.99$) was obtained over 6 orders of magnitude of the standard plasmid DNA concentration (2×10^1 to 2×10^7 copies μL^{-1}), indicating high primer hybridization and extension efficiency.

Anammox and Denitrification Rates. Sediment slurry experiments were conducted to measure rates of anammox and denitrification with a nitrogen isotope tracing method. Detailed information on slurry experiments is also described in refs 23,24. In brief, slurries were prepared with fresh sediment and helium-purged overlying water from each site at a sediment/water volume ratio of 1:5. The resulting slurries were transferred into a series of 12-mL glass vials (Exetainer, Labco, High Wycombe, Buckinghamshire, UK) under a helium atmosphere and pre-incubated for 24 h at near *in situ* temperature to eliminate residual nitrate, nitrite and oxygen. The vials were then divided into three groups, which were spiked (through the septa of each vial) with helium-purged stock solutions of (1) ¹⁵NH₄⁺, (2) ¹⁵NH₄⁺ + ¹⁴NO₃⁻, or (3) ¹⁵NO₃⁻. The final concentration of ¹⁵N in each vial was approximately 100 μ mol L⁻¹. Slurry incubation was blocked by injecting 300 μ l of 50% ZnCl₂ solution into each vial after 8h. The incubation time was determined based on a test experiment that showed a linear increase in N₂ production during an 8-h incubation (Supplementary Fig. 3S). Concentrations of ²⁹N₂ and ³⁰N₂ produced during the incubation were measured by membrane inlet mass spectrometry (MIMS)²². Rates of both anammox and denitrification and their respective contribution to total N₂ production were calculated using the methods given in refs 15, 25.

Dissimilatory Nitrate Reduction to Ammonium (DNRA) Rates. DNRA rates in slurry experiments were measured with a nitrogen isotope tracing method as described in ref. 54. Slurries were made with fresh sediment and helium-purged site water at a sediment-water volume ratio of 1:5 and transferred into 12-mL helium-flushed glass vials (Exetainer, Labco, High Wycombe, Buckinghamshire, UK). Subsequently, they were pre-incubated to eliminate residual nitrate, nitrite, and oxygen at near *in situ* temperature. The slurry vials were then spiked with ¹⁵NO₃⁻ (final concentration ca. 100 μ mol L⁻¹), and one half of the replicates were designated as initial samples and preserved with 300 μ L of 50% ZnCl₂. The remaining slurries were shaken (200 rpm), incubated for approximately 8h at near *in situ* temperature, and preserved as final samples with ZnCl₂ solution, as described for initial samples. The concentrations of ¹⁵NH₄⁺ produced via DNRA in the slurry incubations were analyzed by the Oxidation/MIMS (OX/MIMS) method⁵⁴. Potential DNRA rates were estimated according to the changes in ¹⁵NH₄⁺ concentration during the incubations⁵⁴.

Potential Ammonium Oxidation Rates. Potential nitrification rates were quantified in triplicate according to ref. 58. Briefly, 10g of fresh sediment was added to 250-mL Erlenmeyer flasks containing 100 mL of overlying water from the site. NH_4Cl (for ammonium) and KH_2PO_4 (for phosphate) were amended to the flasks at final concentrations of 300 µmol L⁻¹ and 60µmol L⁻¹, respectively. The suspension was incubated in the dark at near *in situ* temperature with continuous shaking at 120 rpm. During

the incubations, subsamples (2 mL) were harvested at 0, 4, 8, and 12 h and were centrifuged, filtered and frozen immediately for nitrate (plus nitrite) analysis. Potential nitrification rates were estimated based on the linear increase of nitrate concentrations in suspension during incubation⁵⁸.

Statistical Analyses. The Mothur program (version 1.33.0) was used to generate rarefaction curves for the observed unique OTUs and to determine the biodiversity indicators (Shannon and Chao 1) for each constructed gene library. The coverage of each clone library (>90% in this study) was estimated by the percentage of the observed number of OTUs divided by the Chao1 estimate (Supplementary Table S4). The ecological distribution of anammox bacterial communities and their correlations with environmental factors were determined by Weighted UniFrac PCoA analysis and canonical correspondence analysis (CCA), respectively, using CANOCO software⁵⁹. Pearson correlation analyses were conducted with SPSS (version 11.5) to examine correlations among anammox bacterial diversity, abundance and environmental factors. In addition, a one-way ANOVA was performed to compare spatial and seasonal differences in nitrogen transformation rates.

References

- 1. Galloway, J. N. *et al.* Transformation of the nitrogen cycle: Recent trends, questions, and potential solutions. *Science* **320**, 889–892 (2008).
- 2. Gruber, N. & Galloway, J. N. An Earth-system perspective of the global nitrogen cycle. Nature 451, 293-296 (2008).
- 3. Seitzinger, S. P. Nitrogen cycle: Out of reach. Nature 452, 162-163 (2008).
- 4. Davidson, E. A. The contribution of manure and fertilizer nitrogen to atmospheric nitrous oxide since 1860. Nature Geosci. 2, 659-662 (2009).
- Ravishankara, A., Daniel, J. & Portmann, R. Nitrous oxide (N₂O): The dominant ozone-depleting substance emitted in the twenty first century. *Science* 326, 123–125 (2009).
- Bodirsky, B. L. et al. Reactive nitrogen requirements to feed the world in 2050 and potential o mitigate nitrogen pollution. Nat. Commun. 5, 3858, doi: 10.1038/ncomms4858 (2014).
- 7. Seitzinger, S. P. et al. Global river nutrient export: A scenario analysis of past and future trends. Glob. Biogeochem. Cycles 24, GB0A08 (2010).
- 8. Diaz, R. J. & Rosenberg, R. Spreading dead zones and consequences for marine ecosystem. Science 321, 926-929 (2008).
- 9. Canfield, D. E., Glazer, A. N. & Falkowski, P. G. The evolution and future of earth's nitrogen cycle. *Science* **330**, 192–196 (2010). 10. Deegan, L. A. *et al.* Coastal eutrophication as a driver of salt marsh loss. *Nature* **490**, 388–392 (2012).
- Cui, S. H., Shi, Y. L., Groffman, P. M., Schlesinger, W. H. & Zhu, Y. G. Centennial-scale analysis of the creation and fate of reactive nitrogen in China (1910–2010). Proc. Natl. Acad. Sci. USA 110, 2052–2057 (2013).
- 12. Ti, C., Pan, J., Xia, X. & Yan, X. A nitrogen budget of mainland China with spatial and temporal variation. *Biogeochemistry* 108, 381-394 (2012).
- 13. Kuypers, M. M. M. et al. Anaerobic ammonium oxidation by anammox bacteria in the Black Sea. Nature 422, 608-611 (2003).
- Mulder, A., Graaf, A. A., Robertson, L. A. & Kuenen, J. G. Anaerobic ammonium oxidation discovered in a denitrifying fluidized bed reactor. FEMS Microbiol. Ecol. 16, 177–184 (1995).
- Thamdrup, B. & Dalsgaard, T. Production of N₂ through anaerobic ammonium oxidation coupled to nitrate reduction in marine sediments. Appl. Environ. Microbiol. 68, 1312–1318 (2002).
- 16. Lam, P. *et al.* Revising the nitrogen cycle in the Peruvian oxygen minimum zone. *Proc. Natl. Acad. Sci. USA* **106**, 4752–4757 (2009). 17. Trimmer, M. & Nicholls, J. C. Production of nitrogen gas via anammox and denitrification in intact sediment cores along a
- continental shelf to slope transect in the North Atlantic. *Limnol. Oceanogr.* 54, 577–589 (2009).
- Jensen, M. M., Lam, P. & Revsbech, N. P. Intensive nitrogen loss over the Omani shelf due to anammox coupled with dissimilatory nitrite reduction to ammonium. *ISME J.* 5, 1660–1670 (2011).
- 19. Lisa, J., Song, B., Tobias, C. R. & Duernberger, K. A. Impacts of freshwater flushing on anammox community structure and activities in the New River Estuary, USA. *Aquat. Microb. Ecol.* **72**, 17–31 (2014).
- 20. Naeher, S. *et al.* Molecular and geochemical constraints on anaerobic ammonium oxidation (anammox) in a riparian zone of the Seine Estuary (France). *Biogeochemistry* **123**, 237–250 (2015).
- 21. Wang, S., Zhu, G. B., Peng, Y. Z., Jetten, M. S. M. & Yin, C. Anammox bacterial abundance, activity, and contribution in riparian sediments of the Pear River Estuary. *Environ. Sci. Technol.* 46, 8834–8842 (2012).
- 22. Hou, L. J. et al. Anaerobic ammonium oxidation (anammox) bacterial diversity, abundance, and activity in marsh sediments of the Yangtze Estuary. J. Geophys. Res. 118, 1237–1246 (2013).
- 23. Risgaard-Petersen, N. et al. Anaerobic ammonia oxidation in an estuarine sediment. Aquat. Microbial. Ecol. 36, 293-304 (2004).
- Engström, P., Dalsgaard, T., Hulth, S. & Aller, R. C. Anaerobic ammonium oxidation by nitrite (anammox): Implications for N₂ production in coastal marine sediments. *Geochim. Cosmochim. Acta* 69, 2057–2065 (2005).
- Trimmer, M., Nicholls, J. C. & Deflandre, B. Anaerobic ammonium oxidation measured in sediments along the Thames Estuary, United Kingdom. Appl. Environ. Microbiol. 69, 6447–6454 (2003).
- Jetten, M. S. M., Op den Camp, H. J. M., Kuenen, J. G. & Strous, M. Description of the order Brocadiales. In: Krieg, N. R. et al. Bergey's Manual of Systematic Bacteriology. Springer, Heidelberg, Germany. pp 506–603 (2010).
- 27. Hu, B. L. et al. Distribution and diversity of anaerobic ammonium-oxidizing bacteria in the sediments of the Qiantang River. Environ. Microbiol. Rep. 4, 540–547 (2012).
- Schmid, M. C. et al. Anaerobic ammonium-oxidizing bacteria in marine environments: Widespread occurrence but low diversity. Environ. Microbiol. 9, 1476–1484 (2007).
- 29. Zhang, Y. et al. Diversity and abundance of aerobic and anaerobic ammonium-oxidizing bacteria in freshwater sediments of the Xinyi River (China). Environ. Microbiol. 9, 2375–2382 (2007).
- Hamersley, M. R. et al. Water column anammox and denitrification in a temperate permanently stratified lake (Lake Rassnitzer, Germany). Systematic Appl. Microbiol. 32, 571–582 (2009).
- Dale, O. R., Tobias, C. R. & Song, B. Biogeographical distribution of diverse anaerobic ammonium oxidizing (anammox) bacteria in Cape Fear River Estuary. *Environ. Microbiol.* 11, 1194–1207 (2009).
- 32. Hong, Y. G., Xu, X. R., Kan, J. J. & Chen, F. Linking seasonal inorganic nitrogen shift to the dynamics of microbial communities in the Chesapeake Bay. *Appl. Microbiol. Biotechnol.* **98**, 3219–3229 (2014).
- Jetten, M. S. M. et al. Anaerobic ammonium oxidation by marine and freshwater Planctomycete-like bacteria. Appl. Microbiol. Biotechnol. 63, 107–114 (2003).

- Meyer, R. L., Risgaard-Petersen, N. & Allen, D. E. Correlation between anammox activity and microscale distribution of nitrite in a subtropical mangrove sediment. *Appl. Environ. Microb.* 71, 6142–6149 (2005).
- 35. Trimmer, M., Nicholls, J. C., Morley, N., Davies, C. A. & Aldridge, J. Biphasic behavior of anammox regulated by nitrite and nitrate in an estuarine sediment. *Appl. Environ. Microbio.* **71**, 1923–1930 (2005).
- 36. Rattray, J. E. et al. Impact of temperature on ladderane lipid distribution in anammox bacteria. Appl. Environ. Microbiol. 76, 1596–1603 (2010).
- 37. Roberts, K. L., Eate, V. M., Eyre, B. D., Holland, D. P. & Cook, P. L. M. Hypoxic events stimulate nitrogen recycling in a shallow salt-wedge estuary: The Yarra River estuary, Australia. *Limnol. Oceanogr.* 57, 1427–1442 (2012).
- Zhao, Y. *et al.* Seasonal variation and controlling factors of anaerobic ammonium oxidation in freshwater river sediments in the Taihu Lake region of China. *Chemosphere* 93, 2124–2131 (2013).
- 39. Schubert, C. J. et al. Anaerobic ammonium oxidation in a tropical freshwater system (Lake Tanganyika), Environ. Microbiol. 8, 1857–1863 (2006).
- 40. Wenk, C. *et al.* Partitioning between benthic and pelagic nitrate reduction in the Lake Lugano south basin. *Limnol. Oceanogr.* **59**, 1421–1433 (2014).
- 41. Erler, D. V., Eyre, B. D. & Davison, L. The contribution of anammox and denitrification to sediment N₂ production in a surface flow constructed wetland. *Environ. Sci. Technol.* **42**, 9144–9150 (2008).
- 42. Rysgaard, S., Glud, R. N., Risgaard-Petersen, N. & Dalsgaard, T. Denitrification and anammox activity in Arctic marine sediments. *Limnol. Oceanogr.* 49, 1493–1502 (2004).
- Hietanen, S. & Kuparinen, J. Seasonal and short-term variation in denitrification and anammox at a coastal station on the Gulf of Finland, Baltic Sea. Hydrobiologia 596, 67–77 (2008).
- 44. Rich, J. J., Dale, O. R., Song, B. & Ward, B. B. Anaerobic ammonium oxidation (anammox) in Chesapeake Bay sediments. *Microb. Ecol.* 55, 311–320 (2008).
- Dong, L. F. *et al.* Changes in benthic denitrification, nitrate ammonification, and anammox process rates and nitrate and nitrite reductase gene abundances along an estuarine nutrient gradient (the Colne Estuary, United Kingdom). *Appl. Environ. Microbiol.* 75, 3171–3179 (2009).
- 46. Koop-Jakobsen, K. & Giblin, A. E. Anammox in tidal marsh sediments: The role of salinity, nitrogen loading, and marsh vegetation. Estuar. Coast. 32, 238-245 (2009).
- 47. Brin, L., Giblin, A. E. & Rich, J. J. Environmental controls of anammox and denitrification in southern New England estuarine and shelf sediments. *Limnol. Oceanogr.* 59, 851–860 (2014).
- 48. Crowe, S. A., Canfield, D. E., Mucci, A., Sundby, B. & Maranger, R. Anammox, denitrification and fixed-nitrogen removal in sediments from the Lower St. Lawrence Estuary. *Biogeosciences* 9, 4309–4321 (2012).
- Bernard, R., Mortazavi, B. & Kleinhuizen, A. A. Dissimilatory nitrate reduction to ammonium (DNRA) seasonally dominates NO₃⁻ reduction pathways in an anthropogenically impacted sub-tropical coastal lagoon. *Biogeochemistry* 125, 47–64 (2015).
- 50. Cao, W. Z. & Wong, M. H. Current status of coastal zone issues and managements in China: A review. *Environ. Int.* 33, 985–992 (2007).
- 51. He, Q. et al. Economic development and coastal ecosystem change in China. Sci. Rep. 4, 5995, doi: 10.1038/srep05995 (2014).
- 52. Sun, Z. et al. China's coastal wetlands: Conservation history, implementation efforts, existing issues and strategies for future improvement. Environ. Int. 79, 25-41 (2015).
- 53. An, S. Q. et al. China's natural wetlands: Past problems, current status, and future challenges. Ambio 36, 335-342 (2007).
- 54. Yin, G. Y., Hou, L. J., Liu, M., Liu, Z. F. & Gardner, W. S. A novel membrane inlet mass spectrometer method to measure ¹⁵NH₄⁺ for isotope enrichment experiments in aquatic ecosystems. *Environ. Sci. Technol.* **48**, 9555–9562 (2014).
- 55. Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. & Higgins, D. G. The CLUSTAL_X windows interface: Flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* 25, 4876–4882 (1997).
- 56. Kumar, S., Tamura, K. & Nei, M. MEGA3: Integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief Bioinform.* 5, 150–163 (2004).
- Tamura, K., Dudley, J., Nei, M. & Kumar, S. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol. Biol. Evol.* 24, 1596–1599 (2007).
- Bernhard, A. E., Tucker, J., Giblin, A. E. & Stahl, D. A. Functionally distinct communities of ammonia-oxidizing bacteria along an estuarine salinity gradient. *Environ. Microbiol.* 9, 1439–1447 (2007).
- ter Braak, C. J. F. & Smilauer, P. CANOCO Reference Manual and CanoDraw for Windows User's Guide: Software for Canonical Community Ordination (version 4.5) Microcomputer Power (Ithaca NY, USA) (2002).

Acknowledgements

This work is funded by the National Natural Science Foundations of China (Nos. 41322002, 41130525, 41071135 and 41271114) and Program for New Century Excellent Talents in University (NCET). We thank Ye Li, Yankun Liu, Yingpeng Yu, Haiying Yue, and Liufu Zeng for collection of sediment samples, and Wayne S. Gardner for discussion.

Author Contributions

L.J.H., Y.L.Z. and M.L. designed the study and wrote the manuscript. L.J.H., Y.L.Z., X.F.L., X.B.L. and J.G. carried out chemical and molecular analyses. G.Y.Y., F.Y.D., F.C. and X.F.J. made great contributions by commenting upon and revising the manuscript.

Additional Information

Supplementary information accompanies this paper at http://www.nature.com/srep

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Hou, L. et al. Anaerobic ammonium oxidation and its contribution to nitrogen removal in China's coastal wetlands. Sci. Rep. 5, 15621; doi: 10.1038/srep15621 (2015).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/