

PART OF A SPECIAL ISSUE ON COASTAL FLOODING AND STORM RISKS
Wave effects on seedling establishment of three pioneer marsh species: survival, morphology and biomechanics

Haobing Cao^{1,2,3}, Zhenchang Zhu^{2,4}, Rebecca James², Peter M. J. Herman^{3,5}, Liquan Zhang¹, Lin Yuan^{1,6,*} and Tjeerd J. Bouma^{2,7,*}

¹State Key Laboratory of Estuarine and Coastal Research, East China Normal University, 200062 Shanghai, China, ²NIOZ Royal Netherlands Institute for Sea Research, Department of Estuarine and Delta Systems, and Utrecht University, PO Box 140, 4400 AC Yerseke, The Netherlands, ³Hydraulic Engineering Department, Delft University of Technology, Delft, The Netherlands, ⁴Institute of Environmental and Ecological Engineering, Guangdong University of Technology, Guangzhou, China, ⁵Deltares, Delft, The Netherlands, ⁶Institute of Eco-Chongming, East China Normal University, 200062 Shanghai, China and ⁷Faculty of Geosciences, Utrecht University, PO Box 80115, 3508 TC Utrecht, The Netherlands

* For correspondence. E-mail Tjeerd.Bouma@nioz.nl or lyuan@sklec.ecnu.edu.cn

Received: 11 February 2019 Returned for revision: 2 May 2019 Editorial decision: 2 August 2019
Published electronically 25 November 2019

- **Background and Aims** It is important to have an in-depth mechanistic understanding of tidal marsh establishment and dynamics to ensure the long-term persistence of these valuable ecosystems. As wave forcing may be expected to impact seedling establishment, we studied the effect of water-imposed drag forces on seedling survival, morphology and biomechanical properties of three marsh pioneer species that are dominant along the salinity gradient in many areas around the world: *Spartina anglica* (salt to brackish), *Scirpus maritimus* (brackish) and *Phragmites australis* (brackish to fresh).
- **Methods** Using a newly developed plant-shaking mesocosm (PSM) that mimicked water-imposed wave drag forces, the effect of wave stress on seedling survival was examined, together with impacts on morphology and biomechanical properties.
- **Key Results** After 7 weeks of exposure to wave stress, lowered seedling survival and growth for all species was revealed. Wave treatments increased the root/shoot biomass ratio to enhance anchorage and made seedlings more flexible (i.e. reduced flexural rigidity), which might be regarded as a mixed outcome between a stress avoidance and stress tolerance strategy.
- **Conclusions** The different biomechanical responses between the three dominant marsh pioneer species, overall, make them less resistant to external stress. Therefore, our results indicate that the likelihood of marshes becoming established is reduced if wave energy increases. Despite the different biomechanical response of these three pioneer species to waves, the seedlings of all species were found to have low resistance to external stresses.

Keywords: Tidal marshes, establishment, seedling, wave effects, biomechanical response, *Spartina anglica*, *Scirpus maritimus*, *Phragmites australis*.

INTRODUCTION

Tidal marsh ecosystems are increasingly valued for their wave attenuation service (Bouma *et al.*, 2005, 2010) even under extreme storm conditions (Möller *et al.*, 2014). In coping with the growing risk of coastal flooding under accelerating sea-level rise (Craft *et al.*, 2008; Lin *et al.*, 2012; Kirwan and Megonigal, 2013), the conservation and restoration of tidal marshes are now widely appreciated as long-term sustainable protective solutions (Borsje *et al.*, 2011; van Slobbe *et al.*, 2013; Temmerman *et al.*, 2013). Tidal marshes are also valued for many other valuable ecosystem services such as conservation of biodiversity, regulation of nutrients via nutrient cycling, and regulation of climate via organic carbon sequestration (Gedan *et al.*, 2009; Barbier *et al.*, 2011; Burden *et al.*, 2013). Although there is strong evidence that tidal marshes can survive sea-level rise due to sediment-trapping plant-growth feedbacks (Kirwan and Megonigal, 2013; Kirwan *et al.*, 2016), over the last few

decades tidal marshes have rapidly degraded on a global scale (Silliman *et al.*, 2009; Tonelli *et al.*, 2010; Temmerman *et al.*, 2012). The high value of tidal marshes and the loss of such important ecosystems emphasize the needs to conserve and restore them. The latter requires in-depth understanding of both tidal marsh (re)establishment and lateral expansion under external stressors (Fagherazzi *et al.*, 2013; Bouma *et al.*, 2014; van Belzen *et al.*, 2017; and references therein).

In many coastal areas around the world, seedling establishment is of particular importance for the (re)colonization of large bare tidal flats. This may be especially important for those erosional marsh edges where the height of the erosional marsh-cliff may prevent plants that are growing on top of this cliff from expanding clonally onto the lower tidal flat in front of the cliff. This kind of disconnection between marsh and tidal flats may be especially important in macro-tidal systems. In these areas, seedling establishment is regarded as the first critical

threshold that needs to be surpassed to initiate the transition from a bare mud flat to vegetated state (Wang and Temmerman, 2013; Balke et al., 2014; Bouma et al., 2016). Once seedlings have established, further marsh expansion may be expected by lateral clonal growth (van der Wal et al., 2008; Vandenbruwaene et al., 2011). Recent studies have shown that for some species waves may hamper seedling establishment (Silinski et al., 2015, 2016) via wave-induced sediment dynamics (Callaghan et al., 2010; Hu et al., 2015; Bouma et al., 2016). However, the direct effect of waves on newly establishing seedlings remains poorly understood. We still lack insight as to whether this may vary among marsh pioneer species that occur along the salinity gradient of estuaries, where wave exposure may also be expected to decrease due to narrowing of the system.

Wave exposure is known to be of critical importance for understanding long-term marsh development (Callaghan et al., 2010; Fagherazzi et al., 2012; Bouma et al., 2014). For example, recent studies have demonstrated that wave exposure can induce cliff formation at the marsh edge, causing lateral erosion (Callaghan et al., 2010; Marani et al., 2011; Fagherazzi et al., 2013; Wang et al., 2017). In contrast, surprisingly little is known on how wave stress affect tidal marsh plants during the establishment phase. Previous studies indicate that waves can affect individual plants, either directly by imposition of drag/pull on the plants (Coops et al., 1991, 1994, 1996; Coops and Van der Velde, 1996; La Nafie et al., 2012; Silinski et al., 2018), or indirectly via sediment scouring around the stems (Bouma et al., 2009a; Silinski et al., 2015). As yet, the direct mechanisms that limit or allow marsh establishment are only starting to be quantified by experimental studies (Silinski et al., 2015).

Current knowledge gaps on the direct effects of waves on plants may be partly due to methodological limitations. Studies assessing the interaction between waves and vegetation have conventionally used wave flumes of different scales (Bouma et al., 2005, 2009b, 2010; Möller et al., 2014; Silinski et al., 2015, 2018). Whereas wave flumes typically provide a perfect method to mimic hydrodynamic conditions, the high construction and operational costs of such infrastructure makes these experiments generally too expensive to do long-term growth experiments or have high numbers of replicates. Long-term studies are commonly carried out in wave tanks (Coops et al., 1996; Coops and Van der Velde, 1996; La Nafie et al., 2012; Wang et al., 2017), in which hydrodynamic conditions are typically less ideal than in flumes, but due to lower costs some level of replication is possible. In this study, we provide an even more simple approach to assess wave effects on plants by designing a tidal plant-shaking mesocosm (PSM). The PSM mimics wave-induced drag stress on the seedlings, by moving the plants (seedling) back and forth through the water, rather than by moving the water around the plants.

Species-specific insight into the responses of pioneers to enhanced wave climate is important to inform future restoration projects. As wave exposure may be expected to decrease with salinity, due to the typical narrowing of the estuarine system, species with a higher salt tolerance will perhaps be better adapted to wave exposure. The present study aims to extend current knowledge by studying the direct effect of wave stress on the survival, morphology and biomechanical properties of newly establishing seedlings of marsh pioneers, using novel tidal PSM. We compare this for three dominant marsh pioneer

species that occur along the estuaries with decreasing salinity gradient: *Spartina anglica* (salt to brackish), *Scirpus maritimus* (brackish) and *Phragmites australis* (brackish to fresh).

MATERIALS AND METHODS

Plant material

Seedlings of *Sp. anglica*, *Sc. maritimus* and *P. australis* were obtained from seeds that had been collected from the Scheldt estuary (The Netherlands) in November 2016. Seeds were cool stored in a fridge at 4 °C until being germinated in a container with an alternating temperature condition (25 °C during the day and 30 °C at night to speed up germination, practical choice). Seeds with a visible germ were identified as seedlings suitable for transplantation. All seedlings were prepared 1 week before transplantation and stored in an incubator with the same germination conditions as described above. The plants were regarded to be in the seedling stage during the whole period of the experiment, as they remained short compared to fully grown individuals (longest shoots = 17.7 ± 2.4 cm for *Sp. anglica*; 34.9 ± 4.3 cm for *Sc. maritimus*; 44.5 ± 7.2 cm for *P. australis*). For each species, we transplanted 24 seedlings individually to PVC pots (160 mm height \times 110 mm inner diameter) by planting them at 1 cm depth of the sediment. The sediment (with an average D50 of 29.93 μ m) used was macrobenthos-free (by sieving) and selected in a primary test to have negligible scouring under our experimental conditions.

Experimental design

The experiment was carried out using four tidal mesocosms in a climate room (NIOZ Royal Netherlands Institute for Sea Research, Yerseke, The Netherlands), where light was provided with 12 h d⁻¹ (550 μ mol m⁻² s⁻¹ photosynthetically active radiation), and temperature was controlled at 25 °C during the day and 18 °C during night (for more details see references in Cao et al., 2018). In each mesocosm, a semi-diurnal 1.5-h inundation regime was mimicked with a mix of fresh water and Scheldt sea water (which gives a salinity of 12.61 ppt). All pots were equally distributed in the four systems with six seedlings of each species per mesocosm, which gives 12 replicate seedlings per species for each treatment. The water depth of all mesocosms was 21 cm from the sediment top during high tide.

Two of the mesocosms were set as controls (C) and the other two were equipped with plant-shakers to impose continuous drag forces during mimicked high tide. Each PSM consists of a piston-moved cuboid metal frame with a cribriform bottom that can hold many pots with plants. To simulate a wave like stress (W treatment), the plant-shaker was inserted onto the top tank of a tidal mesocosm. The piston was provided with air pressure that was controlled by a timer, to allow the device to move back and forth during tidal inundation. The distance of the motion was 30 cm in both directions, over a period of 2.6 s in the experimental tank, during the semi-diurnal 1.5 h of inundation. In this way, we approximated the drag imposed by regular waves with an average frequency of 0.38 Hz. This is to mimic the measured windy weather wave conditions at tidal marshes

in the Scheldt estuary (this compares with ~0.5 Hz for similar significant wind waves at tidal marsh fronts in the Scheldt estuary field sites, see references in Callaghan *et al.*, 2010; Wang *et al.*, 2017).

Plant morphology and biomechanical properties measuring

The mesocosm experiment lasted for 7 weeks, during which seedling health (survived, toppled or dead) and plant height (soil to leaf top) were surveyed weekly. To compare the overall growth rate of seedlings, the plant height change at each week interval was calculated. To test plant stem flexibility under bending forces, as occurs under wave forcing (Rupprecht *et al.*, 2015), all survived seedlings were carefully cleaned from the sediment at the end of the seventh week.

We measured biomechanical traits by three-point bending tests on the 5-cm-long basal stem fragments from survived seedlings, using a universal testing machine (Instron 5942, Canton, MA, USA). The tests were performed by applying a constant extension rate to the midpoint of each stem sample on two supports of the test machine jaw. The following biomechanical traits that related to bending were calculated:

Young's modulus (E ; kPa), which quantifies the stiffness of the stem and describes how much force is needed to bend a stem (a higher E value indicates a stiffer stem), is calculated as the slope of the stress–strain curve in the elastic deformation region;

the *second moment of area* (I ; m⁴), which quantifies the distribution of material around the axis of bending (increases with diameter), is calculated using the formula for triangular cross-sections: $I = (bh^3)/36$, in which b and h are the base and height of the cross-section;

flexural stiffness (EI , N m⁻²), which was calculated by multiplying E (Young's modulus) and I (the second moment of area) to compare the overall flexural rigidity of the seedlings (a higher EI value indicates less flexibility of individual plants).

The stem was considered to break or fold when it reaches the maximum bending stress.

The *breaking force* (F , N) was then calculated to compare the maximum force that samples can withstand before mechanical failure (a higher F value indicates more resistance to tensile stress).

The breaking force is thus a measurement of the external stress that seedlings can resist during the establishment phase (Rupprecht *et al.*, 2015; Silinski *et al.*, 2015, 2018; Vuik *et al.*, 2018). Due to mortality, we used five replicates for each species per treatment for the biomechanical test. For calculation, all the stem samples were deemed as circular in cross-section because hollow stems had not yet developed for all young seedlings in our experiment. Finally, the dry biomass of survived seedlings was measured after oven-drying at 60 °C for 72 h to compare root/shoot biomass.

Statistics

Seedling survival was analysed with the Kaplan–Meier method applying the log-rank for treatment comparisons, with seedling toppling and death set as hazard events. Seedling

growth rates were calculated based on the average change in plant height after 7 weeks. Two-way ANOVAs were carried out to analyse the effects of wave disturbance and species on the growth rate, root/shoot biomass and biomechanical parameters. All results were tested in SPSS 18.0 software (SPSS, Chicago, IL, USA) with a significance level of 5 %. We tested for normality and homoscedasticity of the data, and these assumptions were passed.

RESULTS

Seedling survival and growth

In the control groups (C), all seedlings survived (100 %) throughout the experiment, regardless of species (Fig. 1). However, when subjected to mimicked wave stress (W), the percentage of survived seedlings decreased over time. Kaplan–Meier tests showed that wave treatment significantly reduced overall seedling survival (Fig. 1, Table 1; $P < 0.05$). Although growth of seedlings was non-linear in our experiment (Fig. 2A), for all three pioneer marsh species, the average growth rate calculated from the change in plant height (soil to leaf top) during the 7 weeks was also significantly lower under wave treatments as compared to the control groups (Fig. 2, Table 2; $P < 0.05$). No significant difference in survival or growth rate was observed between the three pioneer marsh species (Figs 1 and 2). These results show that wave exposure is a common important factor that hampers marsh establishment at the seedling phase.

Seedling biomechanical and morphological traits

When comparing the biomechanical strength of the survived seedlings, two-way ANOVAs shows significant main effects of both wave treatment and species (Table 2). In particular, measurements of Young's modulus of elasticity (E ; kPa) indicate that seedlings exposed to the wave treatment were significantly more flexible (i.e. reduced E) (Fig. 3A, $P = 0.022$ for *Sp. anglica*, $P = 0.014$ for *Sc. maritimus*, $P = 0.003$ for *P. australis*). A less stiff stem material together with a reduced second moment of area (I , Fig. 3B, $P = 0.01$ for *Sp. anglica*, $P = 0.011$ for *Sc. maritimus*, $P = 0.347$ for *P. australis*) led to significantly decreased overall flexural stiffness (EI) values of seedling stems for all three marsh species in response to wave treatments (Fig. 3C, $P = 0.017$ for *Sp. anglica*, $P = 0.002$ for *Sc. maritimus*, $P = 0.003$ for *P. australis*). Consistently, we also observed that the breaking force (F) of all three marsh seedlings showed a significant decrease after wave treatments (Fig. 3D, $P = 0.041$ for *Sp. anglica*, $P = 0.033$ for *Sc. maritimus*, $P = 0.008$ for *P. australis*). Thus, the overall results of the three-point bending test showed that marsh seedlings became more flexible and less resistant to tensile stress after continuous wave exposure due to different material properties as well as reduced stem diameter.

However, in spite of these general biomechanical trends, the pattern of parameters shifted between marsh species. For example, *Sp. anglica* seedlings showed the lowest flexural stiffness (EI , Fig. 3C) and lowest resistance to tensile stress (F , Fig. 3D). Stem diameter (I , Fig. 3B) was highest for *Sc. maritimus*




FIG. 1. Percentage of survived, toppled and dead seedlings of three marshes in the mesocosm experiment (C and W indicate the control and wave treatment groups separately).

TABLE 1. Results of Kaplan–Meier log-rank test on the effects of wave treatments and species on seedling survival during the course of the mesocosm experiment. Both toppling and death of seedlings were set at hazard events during the test

Variable	χ^2	d.f.	P
Wave treatment	25.98	1	<0.001
Species	0.32	2	0.85

seedlings, giving them the highest mechanical resistance (F, Fig. 3D). For *P. australis* seedlings, although stem diameter (I, Fig. 3B) was comparable between the control (C) and wave treatment groups (W), the significant decrease in stem flexibility (E and EI; Fig. 2A, C) nevertheless reduced the external stress that they were able to withstand (Fig. 3D).

Measurements of morphological traits at harvest showed that constant wave exposure also significantly increased the root/shoot biomass ratio of survived seedlings (Fig. 4, Table 2, $P < 0.05$). Interestingly, the root/shoot biomass ratio of *Sp. anglica* seedling was significantly higher than for the other two species after wave stress treatment (Fig. 4, $P = 0.023$ compared to *Sc. maritimus*; $P = 0.016$ compared to *P. australis*).

DISCUSSION

An in-depth understanding of how various physical and biological factors affect thresholds of tidal marsh establishment is essential for enabling restoration and sustainable management of these valuable ecosystems (Moreno-Mateos et al., 2012; Bouma et al., 2014). As yet, mechanistic studies on the direct effect of waves on the probability of establishment and developmental consequences for marsh seedlings are only starting to emerge (see references in Silinski et al., 2015 for the influence of different wave regimes on the survival chances of *Sc. maritimus* from different life stages; see references in Rupprecht et al., 2015 for field observations of biophysical






FIG. 2. Plant height change and average growth rate over 7 weeks of three marsh seedlings (data are means \pm s.e.; C and W indicate control and wave treatment groups separately, * $P < 0.05$).

properties of NW European salt marshes; but also see references in Zhu et al., 2019 for the effects of salinity, inundation and seasonality on the biomechanical properties of marsh vegetation). The present study shows, for the first time, the direct effects of wave treatments on the early seedling establishment of three pioneer marsh species, by using tidal PSMs. We found

TABLE 2. Two-way ANOVAs table of main effects of wave treatment and species and their interactions on the plant traits of seedlings

Response variable	Deviance source	d.f.	Mean square	F	P
Growth rate (cm per week)	Wave treatment	1	55.2	8.84	0.005
	Species	2	17.1	2.73	0.077
	Wave treatment × Species	2	6.9	1.10	0.341
Yong's modulus (kPa)	Wave treatment	1	86.2	19.67	<0.001
	Species	2	107.8	24.59	<0.001
	Wave treatment × Species	2	59.3	13.52	<0.001
Second moment of area (m ⁴)	Wave treatment	1	0.35	10.56	0.004
	Species	2	0.76	22.98	<0.001
	Wave treatment × Species	2	0.33	9.93	0.001
Flexural stiffness (N mm ²)	Wave treatment	1	390 940	32.56	<0.001
	Species	2	176 809	14.73	<0.001
	Wave treatment × Species	2	87 799	7.31	0.003
Breaking force (N)	Wave treatment	1	2.83	12.82	0.002
	Species	2	1.11	5.03	0.015
	Wave treatment × Species	2	0.70	3.17	0.061
Root/shoot biomass	Wave treatment	1	1.85	9.84	0.003
	Species	2	0.33	1.78	0.180
	Wave treatment × Species	2	0.69	3.68	0.033

FIG. 3. Biomechanical traits of survived seedlings showing more flexibility after the continuous wave exposure: (A) Young's modulus; (B) second moment of area; (C) flexural stiffness; and (D) breaking force of three pioneer marsh seedlings (data are means \pm s.e.; C and W indicate control and wave treatment groups separately; * $P < 0.05$, ** $P < 0.01$).FIG. 4. Root/shoot biomass ratio of survived seedlings at harvest showing increased below-ground biomass investment after the continuous wave exposure (data are means \pm s.e.; C and W indicate control groups and wave treatment groups separately; * $P < 0.05$).

that wave treatments directly reduced seedling survival, enhanced the root/shoot biomass ratio to enlarge anchorage, and made seedlings more flexible (reduced flexural rigidity). The different biomechanical responses between the three dominant marsh pioneer species: *Sp. anglica* (salt to brackish), *Sc. maritimus* (brackish) and *P. australis* (brackish to fresh), overall, make them less resistant to external stress.

Effects of wave stress on seedling establishment and growth

Our results clearly show for all three pioneer marsh species that waves are a stress that may create a threshold that directly hampers both seedling establishment and seedling growth. Marsh recovery has been regarded as a critical transition

between two alternative stable states (the bare tidal flat state versus the vegetated state; Wang and Temmerman, 2013; van Belzen et al., 2017). The present results thus underline the requirement for wave-free periods to provide ‘windows of opportunity’ for a marsh to establish (Balke et al., 2014; Bouma et al., 2014). Whereas previous studies related the need for a wave-free period to prevent uprooting due to sediment dynamics (Bouma et al., 2014, 2016; Cao et al., 2018), we show here that there are also direct effects on seedling growth. As the windows of opportunity concept for seedling establishment requires a disturbance-free period for the seedlings to grow large enough to be able to resist hydrodynamic disturbances (e.g. by having longer roots and stronger shoots; Balke et al., 2013, 2014; Hu et al., 2015; and references therein), reduced seedling growth implies that marsh establishment could be much more vulnerable to waves than previously anticipated. Moreover, as ship-generated waves and extreme weather events may be expected to increase in the near future (Silinski et al., 2015), wave-free windows of opportunity will become more rare. In these cases, site-specific conditions will have to be taken into account (e.g. how much wave stress would be tolerable, or how many windows of opportunity are available) for engineering or management measures that focus on creating marshes by reducing waves on the foreshore.

Response of seedlings in biomechanical traits

Plasticity induced by wave stress can be beneficial to enhance the performance of aquatic plants in coping with external forces (Puijalon et al., 2005, 2008, 2011). For example, a significant increase in root/shoot biomass ratio indicates an increased investment of seedlings to below-ground biomass to reinforce anchorage strength: a ‘tolerance strategy’ to resist greater drag force. This is in line with more commonly observed anchorage increase under flow in other aquatic plant species, such as *Luronium natans*, *Mentha aquatica*, *Potamogeton coloratus* and *Sparganium emersum* (Puijalon et al., 2008). Unlike the seedlings in the control group that can tolerate higher breaking force, a decrease in flexural rigidity of seedlings under wave stress enables them to expose less frontal surface areas to incoming waves: an ‘avoidance strategy’ to experience lower drag forces. This supports a well-recognized adaptation of plants to cope with exposure to strong hydrodynamics (see references in Gaylord et al., 2003 for kelp canopies; Bouma et al., 2005 for salt marshes; Bal et al., 2011; Puijalon et al., 2011 for freshwater aquatic plants and La Nafie et al., 2012 for seagrasses). In an earlier study by Silinski et al. (2015), the authors reported that a shift between avoidance and tolerance strategies may occur between life stages within one species. We found that both strategies are, to some extent, adopted particularly during the early seedling phase.

In addition to the above-mentioned similar adjustment in biomechanical traits for all three pioneer marsh seedlings, we also compared interspecific differences in mechanical characteristics, as they may play a role in the shifts of species along estuaries with varying wave exposure (Callaghan et al., 2010; Wang et al., 2017). For example, the significantly higher root/shoot biomass ratio of *Sp. anglica* seedlings under wave

stress can be expected to enable their establishment at higher hydrodynamic conditions (e.g. higher wind exposure or under sea-level rise) than the other two pioneer marsh species. The notably higher stiffness for *P. australis* seedlings in the control groups might be due to nodal stabilization in their stems (Spats et al., 1990). This means that when living in sheltered areas with less oscillating stress, aquatic plants of *P. australis* should be more tolerant to wave attack than the other two species by following a ‘tolerance strategy’. The present results extend current knowledge that salinity is an important factor governing spatial species distribution and plant growth (Pennings et al., 2005; Crain et al., 2008), in showing that waves may reduce seedling survival by around 50 %. It should be noted that in our experiment, the species were not always grown at their predominant salinity. That is, *Sp. anglica* was grown at a relatively low salinity whereas *P. australis* was grown at a relatively high salinity compared to their predominant habitat. The latter may have caused a somewhat higher growth rate for *Sp. anglica* and a somewhat lowered growth rate for *P. australis*, with potential effects on exact survival percentages. The general trend that waves hamper seedling survival for pioneer marsh species remains clear. Moreover, recent year-round field observations by Zhu et al. (2019) showing that salinity did not result in major changes in plant biomechanical properties suggest that the present findings of wave-effects on (biomechanical) plant traits is not strongly affected by using a single salinity.

Our results that *Sp. anglica* seedlings had the least flexural rigidity and resistance to tensile stress (as compared to change in biomechanical properties for the other two species seedlings) seem controversial given that *Sp. anglica* is well known as having wave attenuation value with stiff shoots (Bouma et al., 2005). However, considering that individual seedlings were used in the present study and the high variability in the flexural rigidity of different life cycles of marsh plants (Rupprecht et al., 2015), the stem biomass of dense mature plants canopies may compensate for the stiffness in wave dissipation (Bouma et al., 2005; Möller et al., 2014; Rupprecht et al., 2015). Therefore, the present findings are in agreement with earlier studies that highlight the importance of considering plant size and morphology when determining plant biomechanical characteristics for their wave attenuation service at different locations and times (Niklas, 1992; Rupprecht et al., 2015; Silinski et al., 2018; Zhu et al., 2019).

Implications for tidal marsh restoration

In recent decades, physical constraints have been widely recognized to be the overarching control for marsh establishment (Adam, 2002; Friess et al., 2012). The mechanisms responsible for determining critical a hydrodynamic threshold for marsh establishment and long-term evolution have mainly been elucidated by modelling studies (Mariotti and Fagherazzi, 2010). For example, wave stress and the sediment surface erosion they induce are predicted to be the essential processes behind marsh lateral retreat (Callaghan et al., 2010; Tonelli et al., 2010; Marani et al., 2011; Francalanci et al., 2013). Recent experimental studies have extensively looked at the threshold constraints in terms of sediment dynamics (Balke et al., 2014;

Bouma *et al.*, 2016; Cao *et al.*, 2018) or elevation-related inundation (Wang and Temmerman, 2013; van Belzen *et al.*, 2017). In this regard, our results reveal direct wave-stress effects on seedling survival and growth (both restriction and response) and provide a quantitative support for modelling studies to elucidate hydrodynamically driven tidal marsh dynamics. However, wave characteristics (direction, length, height) can vary strongly in space and time (Nielsen, 2009). Due to the impact of increasing ship traffic or deeper water levels following sea-level rise, we may expect more severe wave forcing in an era of global change (Curtiss *et al.*, 2009; Houser, 2010). To better understanding the effect of wave stress on vegetation, it is thus also important to quantify actual wave forcing in the field, and how this changes over time. Such data can then be used to design process-based studies that investigate the effect of waves with varying energy on marsh expansion both by seedling establishment and rhizome expansion (Bouma *et al.*, 2014, 2016; Silinski *et al.*, 2015, 2018).

Overall, the present study indicates that the likelihood of marsh establishment is reduced if wave energy increases. Despite the different biomechanical response of the three pioneer species studied, the seedlings of all species have low resistance to external stresses. This type of experimental knowledge is essential to understand the role that coastal vegetation plays in mitigating risk and defending coastlines under sea-level rise, and to build more reliable process-based models to predict long-term marsh ecosystem dynamics, and evaluate measures for site-specific management and restoration schemes.

FUNDING

This work was financed by the national key research and development programme of China (Grants: 2017YFC0506002, 2016YFE0133700), the National Natural Science Foundation of China (Grant: 41876093), and the CoE-Rammegors project of the Netherlands. Z.Z. was also supported by a project funded by China Postdoctoral Science Foundation (2019M652825).

ACKNOWLEDGEMENTS

We gratefully acknowledge Bert Sinke and the various technicians of NMF of the NIOZ for their help in designing and building the PSMs. Daniel Blok, Lennart van IJzerloo and Jeroen van Dalen are thanked for their help in setting up the experiment.

LITERATURE CITED

- Adam P. 2002. Saltmarshes in a time of change. *Environmental Conservation* 29: 39–61. doi:10.1017/S0376892902000048
- Barbier EB, Hacker SD, Kennedy C, Koch EW, Stier AC, Silliman BR. 2011. The value of estuarine and coastal ecosystem services. *Ecological Monographs* 81: 169–193. doi:10.1890/10–1510.1
- Bal KD, Bouma TJ, Buis K, *et al.* 2011. Trade-off between drag reduction and light interception of macrophytes: comparing five aquatic plants with contrasting morphology. *Functional Ecology* 25: 1197–1205. doi:10.1111/j.1365-2435.2011.01909.x
- Balke T, Webb EL, van den Elzen E, Galli D, Herman PMJ, Bouma TJ. 2013. Seedling establishment in a dynamic sedimentary environment: a conceptual framework using mangroves. *Journal of Applied Ecology* 50: 740–747.
- Balke T, Herman PMJ, Bouma TJ. 2014. Critical transitions in disturbance-driven ecosystems: identifying windows of opportunity for recovery. *Journal of Ecology* 102: 700–708.
- Borsje BW, van Wesenbeeck BK, Dekker F, *et al.* 2011. How ecological engineering can serve in coastal protection. *Ecological Engineering* 37: 113–122.
- Bouma TJ, De Vries MB, Low E, *et al.* 2005. Trade-offs related to ecosystem engineering: a case study on stiffness of emerging macrophytes. *Ecology* 86: 2187–2199.
- Bouma TJ, Friedrichs M, Klaassen P, *et al.* 2009a. Effects of shoot stiffness, shoot size and current velocity on scouring sediment from around seedlings and propagules. *Marine Ecology Progress Series* 388: 293–297.
- Bouma TJ, Friedrichs M, van Wesenbeeck BK, Temmerman S, Graf G, Herman PMJ. 2009b. Density-dependent linkage of scale-dependent feedbacks: a flume study on the intertidal macrophyte *Spartina anglica*. *Oikos* 118: 260–268.
- Bouma TJ, Vries MBD, Herman PMJ. 2010. Comparing ecosystem engineering efficiency of two plant species with contrasting growth strategies. *Ecology* 91: 2696–2704. doi: 10.1890/09-0690.1
- Bouma TJ, van Belzen J, Balke T, *et al.* 2014. Identifying knowledge gaps hampering application of intertidal habitats in coastal protection: opportunities & steps to take. *Coastal Engineering* 87: 147–157.
- Bouma TJ, van Belzen J, Balke T, *et al.* 2016. Short-term mudflat dynamics drive long-term cyclic salt marsh dynamics. *Limnology and Oceanography* 61: 2261–2275.
- Burden A, Garbitt RA, Evans CD, Jones DL, Cooper DM. 2013. Carbon sequestration and biogeochemical cycling in a saltmarsh subject to coastal managed realignment. *Estuarine Coastal and Shelf Science* 120: 12–20.
- Callaghan DP, Bouma TJ, Klaassen P, van der Wal D, Stive MJF, Herman PMJ. 2010. Hydrodynamic forcing on salt-marsh development: distinguishing the relative importance of waves and tidal flows. *Estuarine, Coastal and Shelf Science* 89: 73–88.
- Cao H, Zhu Z, Balke T, Zhang L, Bouma TJ. 2018. Effects of sediment disturbance regimes on *Spartina* seedling establishment: implication for salt marsh creation and restoration. *Limnology and Oceanography* 63: 647–659.
- Coops H, Van der Velde G. 1996. Effects of waves on helophyte stands: mechanical characteristics of stems of *Phragmites australis* and *Scirpus lacustris*. *Aquatic Botany* 53: 175–185. doi:10.1016/0304-3770(96)01026-1
- Coops H, Boeters R, Smit H. 1991. Direct and indirect effects of wave attack on helophytes. *Aquatic Botany* 41: 333–352. doi:10.1016/0304-3770(91)90052-7
- Coops H, Geilen N, van der Velede G. 1994. Distribution and growth of the helophyte species *Phragmites australis* and *Scirpus lacustris* in water depth gradients in relation to wave exposure. *Aquatic Botany* 48: 273–284. doi: 10.1016/0304-3770(94)90020-5
- Coops H, Geilen N, Verheij H, Boeters R, Van der Velde G. 1996. Interactions between waves, bank erosion and emergent vegetation: an experimental study in a wave tank. *Aquatic Botany* 53: 187–198.
- Craft C, Clough J, Ehman J, *et al.* 2008. Forecasting the effects of accelerated sea-level rise on tidal marsh ecosystem services. *Frontiers in Ecology and the Environment* 7: 73–78.
- Crain CM, Albertson LK, Bertness MD. 2008. Secondary succession dynamics in estuarine marshes across landscape-scale salinity gradients. *Ecology* 89: 2889–2899. doi: 10.1890/03-0745
- Curtiss GM, Osborne PD, Horner-Devine AR. 2009. Seasonal patterns of coarse sediment transport on a mixed sand and gravel beach due to vessel wakes, wind waves, and tidal currents. *Marine Geology* 259: 73–85. doi:10.1016/j.margeo.2008.12.009
- Fagherazzi S, Kirwan ML, Mudd MM, *et al.* 2012. Numerical models of salt marsh evolution: ecological, geomorphic, and climatic factors. *Reviews of Geophysics* 50: RG1002.
- Fagherazzi S, Mariotti G, Wiberg PL, MacGlathery K. 2013. Marsh collapse does not require sea-level rise. *Oceanography* 26: 70–77.
- Francalanci S, Bendoni M, Rinaldi M, Solari L. 2013. Ecomorphodynamic evolution of salt marshes: Experimental observations of bank retreat processes. *Geomorphology* 198: 53–65. doi:10.1016/j.geomorph.2013.04.026
- Friess DA, Krauss KW, Horstman EM, *et al.* 2012. Are all intertidal wetlands naturally created equal? Bottlenecks, thresholds and knowledge gaps to mangrove and saltmarsh ecosystems. *Biological Reviews, Cambridge Philosophical Society* 87: 346–366. doi: 10.1111/j.1469-185X.2011.00198.x

- Gaylord B, Denny MW, Koehl MAR. 2003. Modulation of wave forces on kelp canopies by alongshore currents. *Limnology and Oceanography* **48**: 860–871. doi: 10.4319/lo.2003.48.2.0860
- Gedan KB, Silliman BR, Bertness MD. 2009. Centuries of human-driven change in salt marsh ecosystems. *Annual Review of Marine Science* **1**: 117–41.
- Houser C. 2010. Relative importance of vessel-generated and wind waves to salt marsh erosion in a restricted fetch environment. *Journal of Coastal Research* **262**: 230–240. doi:10.2112/08-1084.1
- Hu Z, van Belzen J, van der Wal D, et al. 2015. Windows of opportunity for salt marsh vegetation establishment on bare tidal flats: the importance of temporal and spatial variability in hydrodynamic forcing. *Journal of Geophysical Research: Biogeosciences* **120**: 1450–1469.
- Kirwan ML, Megonigal JP. 2013. Tidal wetland stability in the face of human impacts and sea-level rise. *Nature* **504**: 53–60. doi:10.1038/nature12856
- Kirwan ML, Temmerman S, Skeehean EE, Guntenspergen GR, Fagherazzi S. 2016. Overestimation of marsh vulnerability to sea level rise. *Nature Climate Change* **6**: 253–260.
- La Nafie YA, de los Santos CB, Brun FG, van Katwijk MM, Bouma TJ. 2012. Waves and high nutrient loads jointly decrease survival and separately affect morphological and biomechanical properties in the seagrass *Zostera noltii*. *Limnology and Oceanography* **57**: 1664–1672.
- Lin N, Emanuel K, Oppenheimer M, Vanmarcke E. 2012. Physically based assessment of hurricane surge threat under climate change. *Nature Climate Change* **2**: 462–467. doi:10.1038/nclimate1389
- Marani M, D'Alpaos A, Lanzoni S, Santalucia M. 2011. Understanding and predicting wave erosion of marsh edges. *Geophysical Research Letters* **38**: L21401. 10.1029/2011GL048995
- Mariotti G, Fagherazzi S. 2010. A numerical model for the coupled long-term evolution of salt marshes and tidal flats. *Journal Geophysical Research* **115**: F01004. doi:10.1029/2009JF001326
- Moreno-Mateos D, Power ME, Comin FA, Yockteng R. 2012. Structural and functional loss in restored wetland ecosystems. *PLoS Biology* **10**: e1001247.
- Möller I, Kudella M, Rupprecht F, et al. 2014. Wave attenuation over coastal salt marshes under storm surge conditions. *Nature Geoscience* **7**: 727–731. doi:10.1038/ngeo2251
- Niklas KJ. 1992. *Plant biomechanics*. Chicago: University of Chicago Press.
- Nielsen P. 2009. How storm size matters for surge height. *Coastal Engineering* **56**: 1002–1004.
- Pennings SC, Grant MB, Bertness MD. 2005. Plant zonation in low-latitude salt marshes: disentangling the roles of flooding, salinity and competition. *Journal of Ecology* **93**: 159–167. doi: 10.1111/j.1365-2745.2004.00959.x
- Puijalon S, Bornette G, Sagnes P. 2005. Adaptations to increasing hydraulic stress: morphology, hydrodynamics and fitness of two higher aquatic plant species. *Journal of Experimental Botany* **56**: 777–786. doi:10.1093/jxb/eri063
- Puijalon S, Bouma TJ, Van Groenendaal J, Bornette G. 2008. Clonal plasticity of aquatic plant species submitted to mechanical stress: escape versus resistance strategy. *Annals of Botany* **102**: 989–996. doi:10.1093/aob/mcn190
- Puijalon S, Bouma TJ, Douady CJ, et al. 2011. Plant resistance to mechanical stress: Evidence of an avoidance tolerance trade-off. *New Phytologist* **191**: 1141–1149. doi: 10.1111/j.1469-8137.2011.03763.x
- Rupprecht F, Möller I, Evan B, Spencer T, Jensen K. 2015. Biophysical properties of salt marsh canopies — Quantifying plant stem flexibility and above ground biomass. *Coastal Engineering* **100**: 48–57.
- Silinski A, Heuner M, Schoelynck J, et al. 2015. Effects of wind waves versus ship waves on tidal marsh plants: a flume study on different life stages of *Scirpus maritimus*. *PLoS One* **10**: 20150323. doi:10.1371/journal.pone.0118687
- Silinski A, van Belzen J, Fransen E, et al. 2016. Quantifying critical conditions for seaward expansion of tidal marshes: a transplantation experiment. *Estuarine, Coastal and Shelf Science* **169**: 227–237.
- Silinski A, Schoutens K, Puijalon S, et al. 2018. Coping with waves: plasticity in tidal marsh plants as self-adapting coastal ecosystem engineers. *Limnology and Oceanography* **63**: 799–815.
- Silliman BR, Grosholz T, Bertness MD. 2009. *Human impacts on salt marshes: a global perspective*. Berkeley: University of California Press.
- Spats HC, Speck T, Vogellehner D. 1990. Contributions to the biomechanics of plants. II. Stability against local buckling in hollow plant stems. *Plant Biology* **103**: 123–130.
- Temmerman S, Moonen P, Schoelynck J, Govers G, Bouma TJ. 2012. Impact of vegetation die-off on spatial flow patterns over a tidal marsh. *Geophysical Research Letters* **39**: L03406. doi: 10.1029/2011GL050502
- Temmerman S, Meire P, Bouma TJ, Herman PMJ, Ysebaert T, De Vriend HJ. 2013. Ecosystem-based coastal defence in the face of global change. *Nature* **504**: 79–83. doi:10.1038/nature12859
- Tonelli M, Fagherazzi S, Petti M. 2010. Modeling wave impact on salt marsh boundaries. *Journal of Geophysical Research* **115**: C09028. doi:10.1029/2009JC006026
- van Belzen J, van de Koppel J, Kirwan ML, et al. 2017. Vegetation recovery in tidal marshes reveals critical slowing down under increased inundation. *Nature Communications* **8**: 15811. doi:10.1038/ncomms15811
- van der Wal D, Wielemaker-Van den Dool A, Herman PMJ. 2008. Spatial patterns, rates and mechanisms of saltmarsh cycles (Westerschelde, The Netherlands). *Estuarine, Coastal and Shelf Science* **76**: 357–368.
- Vandenbruwaene W, Temmerman S, Bouma TJ, et al. 2011. Flow interaction with dynamic vegetation patches: implications for biogeomorphic evolution of a tidal landscape. *Journal of Geophysical Research: Earth Surface* **116**: F01008.
- van Slobbe E, de Vriend HJ, Aarninkhof S, Lulofs K, de Vries M, Dircke P. 2013. Building with nature: in search of resilient storm surge protection strategies. *Natural Hazards* **65**: 947–966.
- Wang C, Temmerman S. 2013. Does biogeomorphic feedback lead to abrupt shifts between alternative landscape states? An empirical study on intertidal flats and marshes. *Journal of Geophysical Research* **118**: 229–240. doi: 10.1029/2012JF002474
- Vuik V, Suh Heo HY, Zhu Z, Borsje BW, Jonkman SN. 2018. Stem breakage of salt marsh vegetation under wave forcing: a field and model study. *Estuarine, Coastal and Shelf Science* **200**: 41–58. <https://doi.org/10.1016/j.ecss.2017.09.028>
- Wang H, van der Wal D, Li X, et al. 2017. Zooming in and out: scale dependence of extrinsic and intrinsic factors affecting salt marsh erosion. *Journal of Geophysical Research: Earth Surface* **122**: 1455–1470.
- Zhu Z, Yang Z, Bouma TJ. 2020. Biomechanical properties of marsh vegetation in space and time: effects of salinity, inundation and seasonality. *Annals of Botany* **125**: 277–289.